



Supplemental Label



EPA Reg. No. 59639-35
(EXCEPT CALIFORNIA AND NEW YORK)

DANITOL 2.4 EC SPRAY USE ON: BUSHBERRIES, FRUITING VEGETABLES AND PEA (SUCCULENT)

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS.
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL AND BOOKLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

BUSHBERRIES

CROPS	PESTS	PRODUCT RATES FL OZ/A	SPECIAL INSTRUCTIONS
Bushberries: Blueberry highbush lowbush Elderberry Gooseberry Huckleberry Juneberry Lingonberry Salal	Blueberry Maggot Cherry Fruitworm Cranberry Fruitworm Japanese Beetle Obliquebanded Leafroller Plum Curculio	10 2/3 - 16 (0.2 - 0.3 lb ai/A)	Apply with air or ground equipment in adequate water for uniform coverage (3 - 10 gals/A by air or a minimum of 20 gals/A by ground). Make applications beginning when first pest activity is noticed and repeat as needed but not more often than every 14 days. Resistance Management: Do not make more than 2 <i>Danitol</i> 2.4 EC applications per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests
Do not apply within 3 days of harvest. Do not exceed 2 pts (32 fl oz, 0.6 lb ai) total application of <i>Danitol</i> 2.4 EC per acre per season.			
Currant	Cane Borer	10 2/3 - 16 (0.2 - 0.3 lb ai/A)	Apply as a ground application in a minimum of 50 gals. of water per acre. Make applications beginning when first pest activity is noticed and repeat as needed but not more often than every 7 days. Resistance Management: Do not make more than 3 <i>Danitol</i> 2.4 EC applications per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.
Do not apply within 21 days of harvest. Do not exceed 2 2/3 pts (42 2/3 fl oz, 0.8 lb ai) total application of <i>Danitol</i> 2.4 EC per acre per season.			

FRUITING VEGETABLES

CROPS	PESTS	PRODUCT RATE FL OZ/A	SPECIAL INSTRUCTIONS
Fruiting Vegetables: Bell Pepper Chili Pepper Cooking Pepper Eggplant Ground Cherry (<i>Physalis</i> spp.) Pepino Pimento Sweet Pepper Tomatillo	Fruitworm Stinkbug Tobacco Hornworm Twospotted Spider Mite (except CA) Yellowstriped Armyworm	10 2/3 (0.2 lb ai/A)	Apply with ground or air equipment in adequate water for uniform coverage (25 - 120 gals/A by ground or 5 - 10 gals/A by air). Begin applications when first pest activity is noticed and repeat as needed to maintain control but not more often than every 7 days. Mites: Treatment is recommended when mite populations are just beginning to build (less than 5 motiles/leaf). DO NOT tank mix with any copper formulations. <i>Danitol</i> 2.4 EC may be mixed with and/or alternated with commonly used insecticides and miticides to comply with local Insect Pest Management (IPM) programs. <i>Danitol</i> 2.4 EC may be mixed with and/or alternated with commonly used insecticides and miticides to comply with local IPM programs.
Do not apply within 3 days of harvest Do not exceed 2 2/3 pts (42 2/3 fl oz, 0.8 lb ai) total application of <i>Danitol</i> 2.4 EC per acre per season.			

PEA (Succulent)

CROPS	PESTS	PRODUCT RATE FL OZ/A	SPECIAL INSTRUCTIONS
Pea (Succulent including): English Pea Garden Pea Green Pea Pigeon Pea	Thrips Whitefly Stinkbug Climbing Cutworm Looper Twospotted spiderMite	10 2/3 (0.2 lb ai/A)	Apply with air or ground equipment in adequate water for uniform coverage. Begin as a ground application in a minimum of 10 gals. of water per acre or as an aerial application in a minimum of 3 gals. of water per acre. Begin applications when first pest activity is noticed and repeat as needed to maintain control but not more often than every 7 days.
Do not apply within 7 days of harvest. Do not exceed 2 2/3 pts of <i>Danitol</i> 2.4 EC (42 2/3 fl oz, 0.8 lb ai) per acre per season.			

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICATION DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Copyright © 2006 by Valent U.S.A. Corporation

Danitol is a registered trademark of Sumitomo Chemical Company, Ltd.

Knack is a registered trademark of Valent U.S.A. Corporation

Monitor is a registered trademark of Bayer CropScience

Manufactured for
 Valent U.S.A. Corporation
 P. O. Box 8025
 Walnut Creek, CA 94596-8025
 www.valent.com
 Made in U.S.A.

Form 2006-DAN-0012