

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Indar[®] 75WSP

EPA Reg. No. 62719-421

Disease Control in Blueberry

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Indar[®] 75WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Indar 75WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Indar 75WSP.

Directions for Use

Refer to product label for Indar 75WSP for Mixing and Application Instructions.

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
alternaria (<i>Alternaria</i> spp.) anthracnose (<i>Colletotrichum gloeosporioides</i>) leaf spot and blotch (<i>Mycosphaerella</i> spp., <i>Septoria</i> spp.) mummy berry disease (<i>Monilinia vaccinii-corymbosi</i>) phomopsis twig blight and fruit rot (<i>Phomopsis vaccinii</i>) powdery mildew (<i>Microsphaera vaccinii</i>) rusts (<i>Pucciniastrum</i> spp.)	2 (0.094 lb active)	For mummy berry and phomopsis twig blight, begin applications at early green tip and make subsequent applications at intervals of 8 to 14 days. For control of powdery mildew and other diseases listed, begin applications prior to or at onset of disease development and repeat at 8- to 14-day intervals. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> • Do not make more than 4 applications or apply more than 8 oz of Indar 75WSP (0.38 lb active) per acre per year. • Preharvest Interval: Do not apply within 30 days of harvest. • Do not graze livestock in treated fields or offer treated material as a livestock feed item.

Resistance Management

Indar 75WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Indar 75WSP should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

Plantback Intervals

Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.2 to 1 lb active per acre, the PBI must be 210 days.

®Trademark of Dow AgroSciences LLC

R845-052

EPA accepted: 01/19/07

Initial printing.